arXiv:math/0410239v1 [math.DG] 9 Oct 2004

AN ENERGY-THEORETIC APPROACH TO THE HITCHIN-KOBAYASHI CORRESPONDENCE FOR MANIFOLDS, II

TOSHIKI MABUCHI*

Dedicated to Professor Eugenio Calabi on his eightieth birthday

Abstract. Recently, Donaldson proved asymptotic stability for a polarized algebraic manifold M with polarization class admitting a Kähler metric of constant scalar curvature, essentially when the linear algebraic part H of $\mathrm{Aut}^0(M)$ is semisimple. The purpose of this paper is to give a generalization of Donaldson's result to the case where the polarization class admits an extremal Kähler metric, even when H is not semisimple.

1. Introduction

For a connected polarized algebraic manifold (M, L) with an extremal Kähler metric in the polarization class $c_1(L)_{\mathbb{R}}$, we consider the Kodaira embedding

$$\Phi_m = \Phi_{|L^{\otimes m}|} : M \hookrightarrow \mathbb{P}^*(V_m), \qquad m \gg 1,$$

where $\mathbb{P}^*(V_m)$ denotes the set of all hyperplanes in $V_m := H^0(M, \mathcal{O}(L^m))$ through the origin. For the identity component $\mathrm{Aut}^0(M)$ of the group of holomorphic automorphisms of M, let H denote its maximal connected linear algebraic subgroup. Replacing the ample holomorphic line bundle L by some positive integral multiple of L if necessary, we may assume that the natural H-equivariant maps

$$\mathrm{pr}_m : \otimes^m V_1 \to V_m, \qquad m = 1, 2, \ldots,$$

are surjective (cf. [18], Theorem 3), and may further fix an H-linearization of L, i.e., a lift to L of the H-action on M such that H acts on L as bundle isomorphisms covering the H-action on M. In this paper, applying a method in [15], we shall generalize a result in Donaldson [3] about stability to extremal Kähler cases:

Main Theorem. *For a polarized algebraic manifold (M, L) as above with an extremal Kähler metric in the polarization class, there exists an algebraic torus T in H such that the image $\Phi_m(M)$ in $\mathbb{P}^*(V_m)$ is stable relative to T* (cf. Section 2 and [14]) *for $m \gg 1$.*

In particular in [16], by an argument as in [3], an extremal Kähler metric in a fixed integral Kähler class on a projective algebraic manifold M will be shown to be unique up to the action of the group H.

We now consider the set Δ_L of all algebraic tori T in H for which the statement of Main Theorem is valid. Note that, if T' and T'' are algebraic tori in H with $T' \subset T''$ and $T' \in \Delta_L$, then T'' also belongs to Δ_L (cf. [14], Theorem 3.2). Fix once for all an extremal Kähler metric ω_0 in the polarization class in Main Theorem. By a result of Calabi [1], the identity component K of the group of isometries of (M, ω_0) is a maximal compact

---

*Partially supported by JSPS Grant-in-Aid for Scientific Research B No. 13440023.

connected subgroup of H. For the identity component Z of the center of K, we consider the complexification $Z^{\mathbb{C}}$ of Z in H. In order to prove Main Theorem, it suffices to show

Theorem I. *The torus $Z^{\mathbb{C}}$ belongs to Δ_L.*

For the set Δ of all algebraic tori T in $Z^{\mathbb{C}}$, we consider its subset Δ_0 consisting of all irredundant elements in Δ (cf. Definition 2.4), where $Z^{\mathbb{C}}$ always belongs to Δ_0. Furthermore, related to the obstruction as in [13], we shall show that

$$Z^{\mathbb{C}} \in \Delta_1, \tag{1.1}$$

where Δ_1 is the subset of Δ as in Section 2 below, and is obtained from a weighted version of Donaldson's asymptotic expansion approximating balanced metrics. Then the proof of Theorem I is reduced to showing the following:

Theorem II. $\Delta_L \cap \Delta_0 \;=\; \Delta_0 \cap \Delta_1$.

If the scalar curvature of the extremal Kähler metric ω_0 above is constant, and if the obstruction as in [13] vanishes, then we have both $\Delta_1 = \Delta$ and $\{1\} \in \Delta_0$. Hence in this case, Theorem II shows that $\{1\}$ sits in Δ_L. This then proves the main theorem in [15].

One may ask why relative stability in place of ordinary stability has to be considered in our study. The reason why we choose relative stability is because, in general, the obstruction in [13] to asymptotic semistability does not vanish (cf. [17]). Thus, as to the group action on V_m related to stability, we must replace the full special linear group $\mathrm{SL}(V_m)$ of V_m by its subgroup $G_m(T)$ (see (2.3)), where the algebraic torus T in $Z^{\mathbb{C}}$ is chosen in such a way that the obstruction vanishes for the group $G_m(T)$, i.e., $G_m(T)$ fixes $\hat{M}_m$ (cf. Section 2). Note also that $G_m(T)$ is a direct product of special linear groups. To see why we choose such a group $G_m(T)$ in place of $\mathrm{SL}(V_m)$, we compare our stability with that of holomorphic vector bundles. Recall that a holomorphic vector bundle splitting into a direct sum of stable vector bundles often appears in the boundary of a compactified moduli space of stable vector bundles. Similarly for our stability of manifolds, a splitting phenomenon occurs for V_m in (2.2). Roughly speaking, we consider the moduli space of all M's with fixed decomposition data (2.2), where same type of construction of moduli spaces occurs typically for the Hodge decomposition in the variation of Hodge structures.

We now explain the difficulty which we encounter in applying the method of [15]. Such a difficulty comes up when we use the estimate of Phong and Sturm [21]. By applying a stability criterion in [15] of Hilbert-Mumford's type, we write the vector space $\mathfrak{p}_m$ as an orthogonal direct sum

$$\mathfrak{p}_m \;=\; \mathfrak{p}'_m \oplus \mathfrak{p}''_m, \qquad \text{(cf. Section 3)},$$

and then check the stability of $\hat{M}_m$ along the orbits of the one-parameter subgroups in $G_m(T)$ generated by elements of $\mathfrak{p}''_m$. Though $\mathfrak{p}$ and $\mathfrak{p}''_m$ are transversal by the equality $\mathfrak{p}'_m = \mathfrak{p}_m \cap \mathfrak{p}$, we further need the orthogonality of $\mathfrak{p}$ and $\mathfrak{p}''_m$ in order to apply directly the estimate in [21]. Since such an orthogonality does not generally hold, we are in trouble, but still the situation is not so bad (see (3.17), (3.18)), and this overcomes the difficulty.

It is very likely that the set Δ_L has a natural minimal element closely related to the construction of the torus T_0 in Section 2. To see this, let us consider the case where M is an extremal Kähler toric Fano surface polarized by $L = K_M^{-1}$. Then M is possibly a complex projective plane blown up at r points with $r \leq 3$. If $r = 0$ or 3, then M admits a Kähler-Einstein metric, and the explanation following Theorem II above shows that

Δ_L has the unique minimal element $\{1\}\,(=T_0)$. On the other hand, if $r=1$, then T_0 coincides with $Z^{\mathbb{C}}$, and is the one-dimensional torus generated by the extremal Kähler vector field. Hence, in this case, Δ_L has the unique minimal element T_0. Finally for $r=2$, the involutive holkomorphic symmetry of M switching the blown-up points allows us to regard T_0 as the one-dimensional torus generated by the extremal Kähler vector field. It then follows that T_0 again has to be a minimal element of Δ_L.

2. Notation, convention and preliminaries

Throughout this paper, we fix once for all a pair (M, L) of a connected projective algebraic manifold M and an ample holomorphic line bundle L over M as in the introduction. For V_m in the introduction, we put $N_m := \dim_{\mathbb{C}} V_m - 1$, where the positive integer m is such that L^m is very ample. Let n and d be respectively the dimension of M and the degree of the image $M_m := \Phi_m(M)$ in the projective space $\mathbb{P}^*(V_m)$. Fixing an H-linearization of L as in the introduction, we consider the associated representation: $H \to \mathrm{PGL}(V_m)$. Pulling it back by the finite unramified cover: $\mathrm{SL}(V_m) \to \mathrm{PGL}(V_m)$, we obtain an isogeny

$$\iota : \tilde{H} \to H, \tag{2.1}$$

where $\tilde{H}$ is an algebraic subgroup of $\mathrm{SL}(V_m)$. On the other hand, for an algebraic torus T in H, the H-linearization of L naturally induces a faithful representation

$$H \to \mathrm{GL}(V_m),$$

and this gives a T-action on V_m for each m. Then we have a finite subset $\Gamma_m = \{\chi_1, \chi_2, \dots, \chi_{\nu_m}\}$ of the free Abelian group $\mathrm{Hom}(T, \mathbb{C}^*)$ of all characters of T such that the vector space $V_m = H^0(M, \mathcal{O}(L^m))$ is uniquely written as a direct sum

$$V_m \;=\; \bigoplus_{k=1}^{\nu_m} V_T(\chi_k), \tag{2.2}$$

where for each $\chi \in \mathrm{Hom}(T, \mathbb{C}^*)$, we set $V_T(\chi) := \{s \in V_m\,;\, t\cdot s = \chi(t)\, s \text{ for all } t \in T\}$. Define an algebraic subgroup $G_m = G_m(T)$ of $\mathrm{SL}(V_m)$ by

$$G_m \;:=\; \prod_{k=1}^{\nu_m} \mathrm{SL}(V_T(\chi_k)), \tag{2.3}$$

and the associated Lie subalgebra of $\mathrm{sl}(V_m)$ will be denoted by $\mathfrak{g}_m$. Here, G_m and $\mathfrak{g}_m$ possibly depend on the choice of the algebraic torus T, and if necessary, we denote these by $G_m(T)$ and $\mathfrak{g}_m(T)$, respectively. The T-action on V_m is, more precisely, a right action, while the G_m-action on V_m is a left action. Since T is Abelian, this T-action on V_m can be regarded also as a left action. Note that the group G_m acts diagonally on V_m in such a way that, for each k, the k-th factor $\mathrm{SL}(V_T(\chi_k))$ of G_m acts just on the k-th factor $V_T(\chi_k)$ of V_m. We now put

$$W_m := \{S^d(V_m)\}^{\otimes n+1},$$

where $S^d(V_m)$ denotes the d-th symmetric tensor product of V_m. To the image M_m of M, we can associate a nonzero element $\hat{M}_m$ in W_m^* such that the corresponding element $[\hat{M}_m]$ in $\mathbb{P}^*(W_m)$ is the Chow point of the irreducible reduced algebraic cycle M_m on $\mathbb{P}^*(V_m)$. Note that the G_m-action on V_m naturally induces a G_m-action on W_m and also on W_m^*. As in [14], the subvariety M_m of $\mathbb{P}^*(V_m)$ is said to be *stable relative to T* or *semistable* *relative to* T, according as the orbit $G_m \cdot \hat{M}_m$ is closed in W_m^* or the closure of $G_m \cdot \hat{M}_m$ in W_m^* does not contain the origin of W_m^*.

Take a Hermitian metric h_0 for L such that $c_1(L;h_0)$ is the extremal Kähler metric ω_0 in the Main Theorem. Let E be the extremal Kähler vector field for (M,ω_0), and let $\mathfrak{k}$ be the Lie algebra of K. For ω_0 above, we further define $\Delta_{\min}$ as the set of all $T \in \Delta$ for which the statement of Theorem B in [14] is valid. Then, as the procedure in Section 6 of [14] shows, there exists a unique minimal element, denoted by T_0, of $\Delta_{\min}$ such that $\Delta_{\min} = \{\, T \in \Delta\,;\, T_0 \subset T\,\}$. Then by the notation in (3.1) below, T_0 is the closure in $Z^{\mathbb{C}}$ of the complex Lie subgroup generated by the vector fields

$$E,\ F_k, \quad k = 1,2,\dots\ ,$$

which appear in the asymptotic expansion approximating weighted analogues (cf. [14], 2.6) of balanced metrics. For each $T \in \Delta_{\min}$, we put $\tilde{T} := \iota^{-1}(T)$, and let $G'_m(T)$ and $Z'_m(T)$ be the identity components of $G_m(T) \cap \tilde{H}$ and $G_m(T) \cap Z^{\mathbb{C}}$, respectively. Put

$$\Delta_1 := \{\, T \in \Delta_{\min}\,;\, G'_m(T)\cdot \hat{M}_m = \hat{M}_m \,\}.$$

Definition 2.4. For an algebraic torus T in Δ, we say that T is *irredundant*, if $\dim_{\mathbb{C}} K^{\mathbb{C}} = \dim_{\mathbb{C}} G'_m(T) + \dim_{\mathbb{C}} T$, or equivalently if $\dim_{\mathbb{C}} Z^{\mathbb{C}} = \dim_{\mathbb{C}} Z'_m(T) + \dim_{\mathbb{C}} T$. For instance, if $T = Z^{\mathbb{C}}$, then $Z'_m(T) = \{1\}$. Hence, $Z^{\mathbb{C}}$ is irredundant.

For the complexification $K^{\mathbb{C}}$ of K in H, we define $\tilde{K}^{\mathbb{C}} := \iota^{-1}(K^{\mathbb{C}})$ and its subset $\tilde{Z}^{\mathbb{C}} := \iota^{-1}(Z^{\mathbb{C}})$. For the time being, let $T = Z^{\mathbb{C}}$, and we consider the associated set $\Gamma_m = \{\chi_1, \chi_2 \dots, \chi_{\nu_m}\}$ of characters. Since $K^{\mathbb{C}}$ commutes with $Z^{\mathbb{C}}$, we have the inclusion

$$\tilde{K}^{\mathbb{C}} \quad \subset \quad \mathrm{SL}(V_m) \bigcap \prod_{k=1}^{\nu_m} \mathrm{GL}(V_{Z^{\mathbb{C}}}(\chi_k)). \tag{2.5}$$

Recall that the extremal Kähler vector field E belongs to the Lie algebra of T_0. Hence, a theorem of Calabi [1] shows that $G'_m(Z^{\mathbb{C}}) \subset G'_m(T_0) \subset \tilde{K}^{\mathbb{C}}$. Hence,

$$G'_m(Z^{\mathbb{C}})\cdot \tilde{Z}^{\mathbb{C}} \subset \tilde{K}^{\mathbb{C}}. \tag{2.6}$$

Proof of (1.1). To show (1.1), we compare two groups $[\tilde{K}^{\mathbb{C}}, \tilde{K}^{\mathbb{C}}]$ and $G'_m(Z^{\mathbb{C}})$. By (2.5), we obviously have $[\tilde{K}^{\mathbb{C}}, \tilde{K}^{\mathbb{C}}] \subset G'_m(Z^{\mathbb{C}})$. On the other hand,

$$\dim_{\mathbb{C}} [\tilde{K}^{\mathbb{C}}, \tilde{K}^{\mathbb{C}}] \;=\; \dim_{\mathbb{C}} \tilde{K}^{\mathbb{C}} - \dim_{\mathbb{C}} \tilde{Z}^{\mathbb{C}} \;\geq\; \dim_{\mathbb{C}} G'_m(Z^{\mathbb{C}}),$$

where the last inequality follows from (2.6) in view of the fact that the intersection of $G'_m(Z^{\mathbb{C}})$ and $\tilde{Z}^{\mathbb{C}}$ is a finite group. Now, we see that $G'_m(Z^{\mathbb{C}})$ coincides with $[\tilde{K}^{\mathbb{C}}, \tilde{K}^{\mathbb{C}}]$. Hence $G'_m(Z^{\mathbb{C}})\cdot \hat{M}_m = \hat{M}_m$. Then by $T_0 \subset Z^{\mathbb{C}}$, we now obtain $Z^{\mathbb{C}} \in \Delta_1$, as required. □

Let h be a Hermitian metric for L such that $\omega := c_1(L;h)$ is a K-invariant Kähler metric on M. Define a Hermitian metric ρ_h on V_m by

$$\rho_h(s,s') \;:=\; \int_M (s,s')_{h^m}\, \omega^n, \qquad s, s' \in V_m, \tag{2.7}$$

where $(s,s')_{h^m}$ denotes the function on M obtained as the the pointwise inner product of s, s' by h^m. Let $\mathcal{S} := \{s_0, s_1, \dots, s_{N_m}\}$ be an orthonormal basis for V_m satisfying

$$\rho_h(s_i, s_j) \;=\; \delta_{ij}.$$

Let $T \in \Delta_0 \cap \Delta_1$. Then we say that $\mathcal{S}$ is T-*admissible*, if each $V_T(\chi_k)$, $k = 1, 2, \dots,$ admits a basis $\{s_{k,i}\,;\, i = 1, 2, \dots, n_k\}$ such that

$$s_{l(k,i)} = s_{k,i}, \qquad i = 1, 2, \dots, n_k; \;\; k = 1, 2, \dots, \nu_m, \tag{2.8}$$

where $n_k := \dim_{\mathbb{C}} V_T(\chi_k)$, and $l(k,i) := (i-1) + \Sigma_{k'=1}^{k-1} n_{k'}$ for all k and i (cf. [14]). Let $\mathfrak{t}_c := \mathrm{Lie}(T_c)$ denote the Lie algebra of the maximal compact subgroup T_c of T. Put $q := 1/m$ and $\mathfrak{t}_{\mathbb{R}} := \sqrt{-1}\,\mathfrak{t}_c$. For each $F \in \mathfrak{t}_{\mathbb{R}}$, we define

$$B_q(\omega, F) \;:=\; \frac{n!}{m^n} \sum_{k=1}^{\nu_m} \sum_{i=1}^{n_k} \; e^{-d\chi_k(F)}\, |s_{k,i}|^2_{h^m}, \tag{2.9}$$

where $|s|^2_{h^m} := (s,s)_{h^m}$ for all $s \in V_m$, and $d\chi_k : \mathfrak{t}_{\mathbb{R}} \to \mathbb{R}$ denotes the restriction to $\mathfrak{t}_{\mathbb{R}}$ of the differential at $t = 1$ for the character $\chi_k \in \mathrm{Hom}(T, \mathbb{C}^*)$.

As a final remark in this section, we give an upper bound for degrees of the characters in Γ_m. Let T be an algebraic torus sitting in $Z^{\mathbb{C}}$. By setting $r := \dim_{\mathbb{C}} T$, we identify T with the multiplicative group $(\mathbb{C}^*)^r := \{t = (t_1, t_2, \dots, t_r)\,;\, t_j \in \mathbb{C}^* \text{ for all } j\}$. Since each χ_k in (2.2) may depend on m, the character χ_k will be rewritten as $\chi_{m;k}$ until the end of this section. Then for each $k \in \{1, 2, \dots, \nu_m\}$,

$$\chi_{m;k}(t) \;=\; \prod_{i=1}^{r} t_i^{\alpha(m,k,i)}, \qquad t = (t_1, t_2, \dots . t_r) \in T,$$

for some integers $\alpha(m,k,i)$ independent of the choice of t. Define a nonnegative integer α_m by $\alpha_m := \sup_{k=1}^{\nu_m} \Sigma_{i=1}^r |\alpha(m,k,i)|$. Then we have the following upper bound for α_m:

Lemma 2.10. *For all positive integer m, the inequality $\alpha_m \leq m\,\alpha_1$ holds.*

Proof: Put $S := \mathrm{Ker}\,\mathrm{pr}_m$. Since the subspace S of $\otimes^m V_1$ is preserved by the T-action, we have a T-invariant subspace, denoted by $S^{\perp}$, of $\otimes^m V_1$ such that the vector space $\otimes^m V_1$ is written as a direct sum

$$\otimes^m V_1 \;=\; S \oplus S^{\perp}.$$

Then the restriction of pr_m to $S^{\perp}$ defines a T-equivariant isomorphism $S^{\perp} \cong V_m$. On the other hand, the characters of T appearing in the T-action on $\otimes^m V_1$ are

$$\chi_{\vec{k}}(t) \;:=\; t_1^{\Sigma_{j=1}^m \alpha(1,k_j,1)} t_2^{\Sigma_{j=1}^m \alpha(1,k_j,2)} \cdots t_r^{\Sigma_{j=1}^m \alpha(1,k_j,r)}, \;\; \vec{k} = (k_1, k_2, \dots, k_m) \in I^m,$$

where I^m is the Cartesian product of m-pieces of $I := \{1, 2, \dots, \nu_m\}$. Since $S^{\perp}\,(\cong V_m)$ is a subspace of $\otimes^m V_1$, we now obtain

$$\alpha_m \;\leq\; \max_{\vec{k} \in I^m} \Sigma_{i=1}^r |\Sigma_{j=1}^m \,\alpha(1,k_j,i)| \;\leq\; \max_{\vec{k} \in I^m} \Sigma_{j=1}^m \Sigma_{i=1}^r |\alpha(1,k_j,i)| \;\leq\; m\,\alpha_1,$$

as required. □

Remark 2.11. By the definition of $F(\ell)$ in (3.1) below, Lemma 2.10 above implies that $|d\chi_{m;k}(F(\ell))| \leq C\alpha_1 q$ for some positive real constant C independent of the choice of m and k. Hence in (3.1) below, for each fixed nonnegative integer ℓ, there exists a positive constant C' independent of m and k such that

$$|e^{-d\chi_k(F(\ell))} - 1| \;\leq\; C'q, \qquad k = 1, 2, \dots, \nu_m.$$

In particular, by the notation in (3.2) below, the integral $\int_M \|\tilde{s}_{k,i}\|^2_{h^m} \omega(\ell)^n \,(= e^{-d\chi_k(F(\ell))})$ converges to 1, uniformly in k, as $m \to \infty$.

3. PROOF OF THEOREM II

Fix an arbitrary element T of $\Delta_0 \cap \Delta_1$. Let $m \gg 1$. Then by [14], Theorem B, there exist $F_k \in \mathfrak{t}_{\mathbb{R}}$, real numbers $\alpha_k \in \mathbb{R}$, and smooth real-valued K-invariant functions φ_k, $k = 1, 2, \dots$, on M such that, for each $\ell \in \mathbb{Z}_{\geq 0}$, we have

$$B_q(\omega(\ell), F(\ell)) \;=\; C_{q,\ell} + 0(q^{\ell+2}), \qquad m \gg 1, \tag{3.1}$$

where $F(\ell) := (\sqrt{-1}\,E/2)\,q^2 + \Sigma_{j=1}^{\ell}\, q^{j+2} F_j$, $h(\ell) := h_0 \exp(-\Sigma_{k=1}^{\ell} q^j \varphi_j)$, $C_{q,\ell} := 1 + \Sigma_{j=0}^{\ell}\,\alpha_j q^{j+1}$, and $\omega(\ell) := c_1(L; h(\ell))$. Let us now fix an arbitrary positive integer ℓ. To each T-admissible orthonormal basis $\mathcal{S} := \{s_0, s_1, \dots, s_{N_m}\}$ for $(V_m; \rho_{h(\ell)})$, we associate a basis $\tilde{\mathcal{S}} := \{\tilde{s}_0, \tilde{s}_1, \dots, \tilde{s}_{N_m}\}$ for V_m by

$$\tilde{s}_{k,i} \;=\; e^{-d\chi_k(F(\ell))/2} s_{k,i}, \qquad i = 1, 2, \dots, n_k;\;\; k = 1, 2, \dots, \nu_m, \tag{3.2}$$

where we put $s_{l(k,i)} = s_{k,i}$ and $\tilde{s}_{l(k,i)} = \tilde{s}_{k,i}$ by using the notation in (2.8). We now consider the Kodaira embedding $\Phi_m : M \to \mathbb{P}^*(V_m)$ defined by

$$\Phi_m(x) \;:=\; (\tilde{s}_0(x) : \tilde{s}_1(x) : \cdots : \tilde{s}_{N_m}(x)), \qquad x \in M,$$

where $\mathbb{P}^*(V_m)$ is identified with $\mathbb{P}^{N_m}(\mathbb{C}) = \{\,(z_0 : z_1 : \cdots : z_{N_m})\,\}$ by the basis $\tilde{\mathcal{S}}$. Put $M_m := \Phi_m(M)$. Since $\Delta_L \cap \Delta_0$ is a subset of $\Delta_1 \cap \Delta_0$ (cf. [13], Section 3), the proof of Theorem II (and Main Theorem also) is reduced to showing the following assertion:

Assertion: *The orbit $G_m(T) \cdot \hat{M}_m$ is closed in W_m^*.*

In the Hermitian vector space $(V_m; \rho_{h(\ell)})$, the subspaces $V_T(\chi_k)$, $k = 1, 2, \dots, \nu_m$, are mutually orthogonal. Put

$$K_m \;:=\; \prod_{k=1}^{\nu_m} \mathrm{SU}(V_T(\chi_k); \rho_{h(\ell)}), \qquad \mathfrak{k}_m \;:=\; \bigoplus_{k=1}^{\nu_m} \mathfrak{su}(V_T(\chi_k); \rho_{h(\ell)}).$$

Since T belongs to Δ_1, the group $G'_m = G'_m(T)$ coincides with the isotropy subgroup of G_m at $\hat{M}_m \in W_m^*$. Consider the Lie algebra $\mathfrak{g}'_m := \mathrm{Lie}(G'_m)$ of G'_m. Put $\mathfrak{h} := \mathrm{Lie}(H) = \mathrm{Lie}(\tilde{H})$. Then by $G'_m \subset \tilde{H} \subset \mathrm{SL}(V_m)$, we have the inclusions

$$\mathfrak{g}'_m \;\hookrightarrow\; \mathfrak{h} \;\hookrightarrow\; \mathfrak{sl}(V_m).$$

Put $\mathfrak{k}'_m := \mathrm{Lie}(K'_m)$, where K'_m is the isotropy subgroup of K_m at the point $\hat{M}_m \in W_m^*$. Then $\mathfrak{g}_m$ and $\mathfrak{g}'_m$ are the complexifications of $\mathfrak{k}_m$ and $\mathfrak{k}'_m$, respectively (cf. [1]). Put $\mathfrak{p}_m := \sqrt{-1}\,\mathfrak{k}_m$ and $\mathfrak{p}'_m := \sqrt{-1}\,\mathfrak{k}'_m$. We further define

$$\mathfrak{K}_m \;:=\; \left\{ \bigoplus_{k=1}^{\nu_m} \mathfrak{u}(V_T(\chi_k); \rho_{h(\ell)}) \right\} \bigcap \mathfrak{su}(V_m; \rho_{h(\ell)}), \qquad \mathfrak{P}_m \;:=\; \sqrt{-1}\,\mathfrak{K}_m.$$

By the above inclusions of Lie algebras (see also (2.5)), we can regard $\mathfrak{p} := \sqrt{-1}\,\mathfrak{k}$ as a Lie subalgebra of $\mathfrak{P}_m$. Let ω_{FS} be the Fubini-Study metric on $\mathbb{P}^*(V_m)$ defined by

$$\omega_{\mathrm{FS}} \;:=\; (\sqrt{-1}/2\pi)\,\partial\bar{\partial} \log(\Sigma_{\alpha=0}^{N_m} |z_\alpha|^2).$$

For each $Q \in \mathfrak{P}_m$, let $\mathcal{Q}$ be the associated holomorphic vector field on $\mathbb{P}^*(V_m)$. By the notation for $t = 0$ in Step 1 later in Proof of Assertion, we obtain a vector field $\mathcal{Q}_{TM_m}$ on

M_m via the orthogonal projection of $\mathcal{Q}$ along M_m to tangential directions. Then we have a unique real-valued function φ_Q on $\mathbb{P}^*(V_m)$ satisfying both $\int_{\mathbb{P}^*(V_m)} \varphi_Q\, \omega_{\mathrm{FS}}^{N_m} = 0$ and

$$i_{\mathcal{Q}}(\omega_{\mathrm{FS}}/m) = (\sqrt{-1}/2\pi)\, \bar{\partial}\varphi_Q.$$

Let $\square_{M,\mathrm{FS}} := -\,\bar{\partial}^*\bar{\partial}$ denote the Laplacian on functions on the Kähler manifold $(M, \Phi_m^*\omega_{\mathrm{FS}})$. Define a positive semidefinite K'_m-invariant inner product $(\;,\;)$ on $\mathfrak{P}_m$ by setting

$$\begin{aligned}
(Q_1, Q_2) \;&:=\; \frac{1}{m^2}\int_{M_m} ((\mathcal{Q}_1)_{TM_m}, (\mathcal{Q}_2)_{TM_m})_{\omega_{\mathrm{FS}}}\, \omega_{\mathrm{FS}}^n \\
&=\; \frac{\sqrt{-1}}{2\pi}\int_{M_m} \partial\varphi_{Q_2}\wedge\bar{\partial}\varphi_{Q_1}\wedge n\,\omega_{\mathrm{FS}}^{n-1} \;=\; \int_{M_m} (\bar{\partial}\varphi_{Q_1}, \bar{\partial}\varphi_{Q_2})_{\omega_{\mathrm{FS}}}\,\omega_{\mathrm{FS}}^n \\
&=\; -\int_M \varphi_{Q_1}(\square_{M,\mathrm{FS}}\varphi_{Q_2})\,\Phi_m^*\omega_{\mathrm{FS}}^n \;\in \mathbb{R}
\end{aligned}$$

for all Q_1, $Q_2 \in \mathfrak{P}_m$. Restrict this inner product to $\mathfrak{p}_m$. Then the inner product $(\;,\;)$ on $\mathfrak{p}_m$ is positive definite on $\mathfrak{p}$ and hence on $\mathfrak{p}'_m$. As vector spaces, $\mathfrak{P}_m$ and $\mathfrak{p}_m$ are written respectively as orthogonal direct sums

$$\mathfrak{P}_m \;=\; \mathfrak{p}\oplus\mathfrak{p}^{\perp}, \qquad \mathfrak{p}_m \;=\; \mathfrak{p}'_m \oplus \mathfrak{p}''_m,$$

where $\mathfrak{p}^{\perp}$ is the orthogonal complement of $\mathfrak{p}$ in $\mathfrak{P}_m$, and moreover $\mathfrak{p}''_m$ is the orthogonal complements of $\mathfrak{p}'_m$ in $\mathfrak{p}_m$ (cf. [15]). Hence if $Q \in \mathfrak{p}^{\perp}$, then for any holomorphic vector field $\mathcal{W}$ on M_m, we have

$$\begin{aligned}
&\frac{1}{m^2}\int_{M_m} (\mathcal{Q}_{TM_m}, \mathcal{W}_{TM_m})_{\omega_{\mathrm{FS}}}\,\omega_{\mathrm{FS}}^n \\
&=\; \int_{M_m} (\bar{\partial}\varphi_Q,\, \bar{\theta}_0 + \bar{\partial}(\varphi_{W^1} + \sqrt{-1}\,\varphi_{W^2}))_{\omega_{\mathrm{FS}}}\,\omega_{\mathrm{FS}}^n \\
&=\; \frac{\sqrt{-1}}{2\pi}\int_{M_m} \big\{\,\theta_0\wedge\bar{\partial}\varphi_Q \;+\; \partial(\varphi_{W^1} - \sqrt{-1}\,\varphi_{W^2})\wedge\bar{\partial}\varphi_Q\big\}\wedge n\,\omega_{\mathrm{FS}}^{n-1} \;=\; 0,
\end{aligned}$$

where $i_{\mathcal{W}}(\omega_{\mathrm{FS}}/m)$ on M_m is known to be expressible as $\bar{\theta}_0 + \bar{\partial}(\varphi_{W^1} + \sqrt{-1}\,\varphi_{W^2})$ for some holomorphic 1-form θ_0 on M_m and elements W^1, W^2 in $\mathfrak{p}$. We consider the open neighbourhood (cf. [15])

$$U_m \;:=\; \{\, X \in \mathfrak{p}''_m\,;\, \zeta(\mathrm{ad}\,X)\mathfrak{p}'_m \cap \mathfrak{p}''_m = \{0\}\,\}$$

of the origin in $\mathfrak{p}''_m$, where $\zeta : \mathbb{R} \to \mathbb{R}$ is a real analytic function defined by $\zeta(x) := x(e^x + e^{-x})/(e^x - e^{-x})$, $x \neq 0$, and $\zeta(0) = 0$. By operating $(\sqrt{-1}/2\pi)\,\partial\bar{\partial}\log$ on both sides of (3.1), we obtain

$$\Phi_m^*\omega_{\mathrm{FS}} \;\equiv\; m\,\omega(\ell), \qquad \mod q^{\ell+2}. \tag{3.3}$$

For an element X of $\mathfrak{P}_m$ (later we further assume $X \in \mathfrak{p}''_m$), there exists a T-admissible orthonormal basis $\mathcal{T} := \{\tau_0, \tau_1, \dots, \tau_{N_m}\}$ for $(V_m, \rho_{h(\ell)})$ such that the infinitesimal action of X on V_m can be diagonalized in the form

$$X\cdot\tau_\alpha \;=\; \gamma_\alpha(X)\,\tau_\alpha$$

for some real constants $\gamma_\alpha = \gamma_\alpha(X)$, $\alpha = 0, 1, \dots, N_m$, satisfying $\Sigma_{\alpha=0}^{N_m}\gamma_\alpha(X) = 0$. As in (3.2), we consider the associated basis $\tilde{\mathcal{T}} = \{\tilde{\tau}_0, \tilde{\tau}_1, \dots, \tilde{\tau}_{N_m}\}$ for V_m, where $\tilde{\tau}_{k,i} :=$

$e^{-d\chi_k(F(\ell))/2}\tau_{k,i}$. By setting

$$\lambda_X(e^t) := \exp(t\,X), \qquad t \in \mathbb{R},$$

we consider the one-parameter group $\lambda_X : \mathbb{R}_+ \to \{\,\Pi_{k=1}^{\nu_m} \mathrm{GL}(V_T(\chi_k))\,\} \cap \mathrm{SL}(V_m)$ associated to X. Then $\lambda_X(e^t)\cdot\tau_\alpha = e^{t\gamma_\alpha}\tau_\alpha$ for all α and all $t \in \mathbb{R}$. Moreover,

$$\Phi_m^*\varphi_X \;=\; \frac{\Sigma_{\alpha=0}^{N_m}\gamma_\alpha(X)\,|\tilde{\tau}_\alpha|^2}{m\,\Sigma_{\alpha=0}^{N_m}|\tilde{\tau}_\alpha|^2}, \qquad X \in \mathfrak{P}_m. \tag{3.4}$$

Let η_m be the Kähler form on M defined by $\eta_m := (1/m)\,\Phi_m^*\omega_{\mathrm{FS}}$. To each $X \in \mathfrak{P}_m$, we can associate a real constant c_X such that $\phi_X := c_X + \Phi_m^*\varphi_X$ on M satisfies

$$\int_M \phi_X\,\eta_m^n \;=\; 0.$$

Proof of Assertion: Fix an arbitrary element $0 \neq X$ of $\mathfrak{p}''_m$, and define a real-valued function $f_{X,m}(t)$ on $\mathbb{R}$ by

$$f_{X,m}(t) \;:=\; \log\|\lambda_X(e^t)\cdot\hat{M}_m\|_{\mathrm{CH}(\rho_{h(\ell)})}.$$

For this X, we consider the associated $\gamma_\alpha(X)$, $\alpha = 0,1,\dots,N_m$, defined in the above. From now on, X regarded as a holomorphic vector field on $\mathbb{P}^*(V_m)$ will be denoted by $\mathcal{X}$. By [26] (see also [14], 4.5), we have $\ddot{f}_{X,m}(t) \geq 0$ for all t. Then by [15], Lemma 3.4, it suffices to show the existence of a real number $t_X^{(m)}$ such that

$$\dot{f}_{X,m}(t_X^{(m)}) = 0 < \ddot{f}_{X,m}(t_X^{(m)}) \quad\text{ and }\quad t_X^{(m)}\cdot X \in U_m. \tag{3.5}$$

In the below, real numbers C_i, $i = 1,2,\dots$, always mean positive real constants independent of the choice of m and X. Moreover by abuse of terminology, we write $m \gg 1$, if m satisfies $m \geq m_0$ for a sufficiently large m_0 independent of the choice of X. Then the proof of Assertion will be divided into the following eight steps:

Step 1. Put $\lambda_t := \lambda_X(e^t)$ and $M_{m,t} := \lambda_t(M_m)$ for each $t \in \mathbb{R}$. Metrically, we identify the normal bundle of $M_{m,t}$ in $\mathbb{P}^*(V_m)$ with the subbundle $TM_{m,t}^\perp$ of $T\mathbb{P}^*(V_m)_{|M_{m,t}}$ obtained as the orthogonal complement of $TM_{m,t}$ in $T\mathbb{P}^*(V_m)_{|M_{m,t}}$. Hence, $T\mathbb{P}^*(V_m)_{|M_{m,t}}$ is differentiably written as the direct sum $TM_{m,t}\oplus TM_{m,t}^\perp$. Associated to this, the restriction $\mathcal{X}_{|M_{m,t}}$ of $\mathcal{X}$ to $M_{m,t}$ is written as

$$\mathcal{X}_{|M_{m,t}} \;=\; \mathcal{X}_{TM_{m,t}} \oplus \mathcal{X}_{TM_{m,t}^\perp}$$

for some smooth sections $\mathcal{X}_{TM_{m,t}}$ and $\mathcal{X}_{TM_{m,t}^\perp}$ of $TM_{m,t}$ and $TM_{m,t}^\perp$, respectively. Then the second derivative $\ddot{f}_{X,m}(t)$ is (see for instance [14], [21]) given by

$$\ddot{f}_{X,m}(t) \;=\; \int_{M_{m,t}} |\mathcal{X}_{TM_{m,t}^\perp}|^2_{\omega_{\mathrm{FS}}}\,\omega_{\mathrm{FS}}^n \;\geq\; 0. \tag{3.6}$$

Since the Kodaira embedding $\Phi^{\tilde{\mathcal{T}}} : M \to \mathbb{P}^{N_m}(\mathbb{C})$ defined by

$$\Phi^{\tilde{\mathcal{T}}}(p) := (\tilde{\tau}_0(p) : \tilde{\tau}_1(p) : \cdots : \tilde{\tau}_{N_m}(p))$$

coincides with Φ_m above up to an isometry of $(V_m, \rho_{h(\ell)})$, we may assume without loss of generality that $\Phi^{\tilde{\mathcal{T}}}$ is chosen as Φ_m.

Step 2. In view of the orthogonal decomposition $\mathfrak{P}_m = \mathfrak{p}^\perp \oplus \mathfrak{p}$, we can express X as an othogonal sum

$$X \;=\; X' + X''$$

for some $X' \in \mathfrak{p}$ and $X'' \in \mathfrak{p}^\perp$. Since $\omega(\ell)$ is K-invariant (cf. [14]), the group K acts isometrically on $(V_m, \rho_{h(\ell)})$. Now, there exists a T-admissible orthonormal basis $\mathcal{B} := \{\beta_0, \beta_1, \dots, \beta_{N_m}\}$ for V_m such that the infinitesimal action of X'' on V_m is written as

$$X'' \cdot \beta_\alpha \;=\; \gamma_\alpha(X'')\, \beta_\alpha, \qquad \alpha = 0, 1, \dots, N_m,$$

for some real constants $\gamma_\alpha(X'')$, $\alpha = 0, 1, \dots, N_m$, satisfying $\Sigma_{\alpha=0}^{N_m} \gamma_\alpha(X'') = 0$. By the notation as in (3.2), we consider the associated basis $\tilde{\mathcal{B}} := \{\tilde\beta_0, \tilde\beta_1, \dots, \tilde\beta_{N_m}\}$ for V_m. Then

$$\phi_{X''} \;=\; \frac{\Sigma_{\alpha=0}^{N_m} \hat\gamma_\alpha(X'')\, |\tilde\beta_\alpha|^2}{m\, \Sigma_{\alpha=0}^{N_m} |\tilde\beta_\alpha|^2}, \tag{3.7}$$

where $\hat\gamma_\alpha(X'') := \gamma_\alpha(X'') + m\, c_{X''}$. Now, X' and X'' regarded as holomorphic vector fields on $\mathbb{P}^*(V_m)$ will be denoted by $\mathcal{X}'$ and $\mathcal{X}''$, respectively. Associated to the expression $T\mathbb{P}^*(V_m)_{|M_{m,t}} = TM_{m,t} \oplus TM_{m,t}^\perp$ as differentiable vector bundles, the restrictions $\mathcal{X}'_{|M_{m,t}}$ $\mathcal{X}''_{|M_{m,t}}$ of $\mathcal{X}'$ and $\mathcal{X}''$ to $M_{m,t}$ are respectively written as

$$\mathcal{X}'_{|M_{m,t}} \;=\; \mathcal{X}'_{TM_{m,t}} \oplus \mathcal{X}'_{TM_{m,t}^\perp} \quad\text{ and }\quad \mathcal{X}''_{|M_{m,t}} \;=\; \mathcal{X}''_{TM_{m,t}} \oplus \mathcal{X}''_{TM_{m,t}^\perp},$$

where $\mathcal{X}'_{TM_{m,t}}$, $\mathcal{X}''_{TM_{m,t}}$ are smooth sections of $TM_{m,t}$, and $\mathcal{X}'_{TM_{m,t}^\perp}$ $\mathcal{X}''_{TM_{m,t}^\perp}$ are smooth sections of $TM_{m,t}^\perp$. Then by $X' \in \mathfrak{p}$, we have

$$\mathcal{X}'_{TM_{m,t}^\perp} \;=\; 0, \quad \text{i.e.,} \quad \mathcal{X}_{TM_{m,t}^\perp} \;=\; \mathcal{X}''_{TM_{m,t}^\perp}. \tag{3.8}$$

Step 3. Since T is irredundant, we have $\mathfrak{g}'_m(T) + \mathfrak{t} = \mathfrak{k}^{\mathbb{C}}$, i.e., $\mathfrak{p}'_m + \sqrt{-1}\,\mathfrak{t}_c = \mathfrak{p}$, where these are equalities as Lie subalgebras of $\mathfrak{h}$. From now on until the end of this step, as in the preceding steps, we regard both $\mathfrak{p}'_m$ and $\mathfrak{k}^{\mathbb{C}}$ as Lie subalgebras of $\mathfrak{sl}(V_m)$. Hence, as Lie subalgebras of $\mathfrak{sl}(V_m)$, we have

$$\mathfrak{p} \;=\; \mathfrak{p}'_m + \sqrt{-1}\,\tilde{\mathfrak{t}}_c.$$

where we put $\tilde{\mathfrak{t}}_c := \mathrm{Lie}(T_c)$ for the maximal compact subgroup $\tilde T_c$ of $\tilde T := \iota^{-1}(T)$. Then we can write $X' \in \mathfrak{p}$ as a sum

$$X' \;=\; Y + W$$

for some $Y \in \mathfrak{p}'_m$ and some $W \in \sqrt{-1}\,\tilde{\mathfrak{t}}_c$. Note that the holomorphic vector fields $\mathcal{Y}$ and $\mathcal{W}$ on $\mathbb{P}^*(V_m)$ induced by Y and W, respectively, are tangent to M. By $[Y, W] = 0$, there exists a T-admissible orthonormal basis $\{\sigma_0, \sigma_1, \dots, \sigma_{N_m}\}$ for V_m such that

$$\begin{cases} Y \cdot \sigma_\alpha \;=\; \gamma_\alpha(Y)\, \sigma_\alpha, & \alpha = 0, 1, \dots, N_m; \\ W \cdot \sigma_{k,i} \;=\; b_k\, \sigma_{k,i}, & k = 1, 2, \dots, \nu_m, \end{cases}$$

for some real constants $\gamma_\alpha(Y)$ and b_k, where in the last equality, $\sigma_{k,i} := \sigma_{l(k,i)}$ as in (2.8). By setting $\tilde\sigma_{k,i} \;=\; e^{-d\chi_k(F(\ell))/2}\sigma_{k,i}$, we later consider the basis $\{\tilde\sigma_0, \tilde\sigma_1, \dots, \tilde\sigma_{N_m}\}$ for V_m. Note that $\Sigma_{\alpha=0}^{N_m} \gamma_\alpha(Y) = \Sigma_{k=1}^{\nu_m} n_k b_k = 0$. Since both X and Y belong to $\mathfrak{p}_m$, it follows from $X = X' + X'' = Y + W + X''$ that

$$\Sigma_{i=1}^{n_k} \gamma_{k,i}(X) \;=\; \Sigma_{i=1}^{n_k} \gamma_{k,i}(Y) \;=\; 0 \quad\text{and}\quad n_k b_k \;=\; -\Sigma_{i=1}^{n_k} \gamma_{k,i}(X'') \quad \text{for all } k,$$

where $\gamma_{k,i} := \gamma_{l(k,i)}$ as in (2.8). Note that $X \in \mathfrak{p}''_m$ and $X'' \in \mathfrak{p}^{\perp}$, where by $\mathfrak{p}'_m + \sqrt{-1}\,\mathfrak{t} = \mathfrak{p}$, the space $\mathfrak{p}^{\perp}$ is perpendicular to $\mathfrak{p}'_m$. Then by $Y \in \mathfrak{p}'_m$ and $X = Y + W + X''$, we have

$$(Y,Y) + (Y,W) \,=\, (Y,X) \,=\, 0$$

in terms of the inner product (,) on $\mathfrak{P}_m$. Hence $(Y,Y) \,=\, -\,(Y,W) \,\le\, \sqrt{(Y,Y)(W,W)}$. It now follows that

$$\int_{M_m} |\mathcal{Y}_{|M_m}|^2_{\omega_{\rm FS}}\,\omega^n_{\rm FS} \,=\, m^2(Y,Y) \,\le\, m^2(W,W) \,=\, \int_{M_m} |\mathcal{W}_{|M_m}|^2_{\omega_{\rm FS}}\omega^n_{\rm FS}. \tag{3.9}$$

The integral on the right-hand side is, for $m \gg 1$,

$$\begin{aligned}
&\int_M \frac{(\Sigma_{k=1}^{\nu_m}\Sigma_{i=1}^{n_k}|\tilde\sigma_{k,i}|^2_{h(\ell)})(\Sigma_{k=1}^{\nu_m}\Sigma_{i=1}^{n_k}b_k^2|\tilde\sigma_{k,i}|^2_{h(\ell)}) - (\Sigma_{k=1}^{\nu_m}\Sigma_{i=1}^{n_k}b_k|\tilde\sigma_{k,i}|^2_{h(\ell)})^2}{(\Sigma_{k=1}^{\nu_m}\Sigma_{i=1}^{n_k}|\tilde\sigma_{k,i}|^2_{h(\ell)})^2} m^n\eta_m^n\\
&\le\; m^n \int_M \frac{\Sigma_{k=1}^{\nu_m}\Sigma_{i=1}^{n_k}b_k^2|\tilde\sigma_{k,i}|^2_{h(\ell)}}{\Sigma_{k=1}^{\nu_m}\Sigma_{i=1}^{n_k}|\tilde\sigma_{k,i}|^2_{h(\ell)}}\,\eta_m^n\\
&\le\; \frac{n!}{2}\int_M \Sigma_{k=1}^{\nu_m}\Sigma_{i=1}^{n_k}b_k^2|\tilde\sigma_{k,i}|^2_{h(\ell)}\,\eta_m^n \;\le\; C_1\Sigma_{k=1}^{\nu_m}n_k b_k^2
\end{aligned}$$

for some C_1, where in the last two inequalities, we used Remark 2.11 in Section 2. Hence, by setting $\hat\gamma_\alpha(Y) := \gamma_\alpha(Y) + mc_Y$, we see from (3.9) that

$$\begin{aligned}
&\int_M \frac{(\Sigma_{\alpha=0}^{N_m}|\tilde\sigma_\alpha|^2_{h(\ell)})(\Sigma_{\alpha=0}^{N_m}\hat\gamma_\alpha(Y)^2|\tilde\sigma_\alpha|^2_{h(\ell)}) \;-\; (\Sigma_{\alpha=0}^{N_m}\hat\gamma_\alpha(Y)|\tilde\sigma_\alpha|^2_{h(\ell)})^2}{(\Sigma_{\alpha=0}^{N_m}|\tilde\sigma_\alpha|^2_{h(\ell)})^2}\,\eta_m^n\\
&\qquad\le\quad q^n C_1\,\Sigma_{k=1}^{\nu_m}\,n_k b_k^2.
\end{aligned} \tag{3.10}$$

Define real numbers f_1 and f_2 by

$$\begin{cases} f_1 & := \int_M(\Sigma_{\alpha=0}^{N_m}|\tilde\sigma_\alpha|^2_{h(\ell)})^{-1}(\Sigma_{\alpha=0}^{N_m}\hat\gamma_\alpha(Y)^2|\tilde\sigma_\alpha|^2_{h(\ell)})\,\eta_m^n,\\ f_2 & := \int_M\{\,(\Sigma_{\alpha=0}^{N_m}|\tilde\sigma_\alpha|^2_{h(\ell)})^{-1}(\Sigma_{\alpha=0}^{N_m}\hat\gamma_\alpha(Y)\,|\tilde\sigma_\alpha|^2_{h(\ell)})\,\}^2\,\eta_m^n. \end{cases}$$

If $f_1 \ge 2f_2$, then by (3.10) and Remark 2.11, we have

$$\Sigma_{\alpha=0}^{N_m}\,\hat\gamma_\alpha(Y)^2 \;\le\; C_2\,\Sigma_{k=1}^{\nu_m}\,n_k b_k^2, \qquad \text{if } m \gg 1,$$

for some C_2. Next, assume $f_1 < 2f_2$. Then for

$$\phi_Y := (m\Sigma_{\alpha=0}^{N_m}|\tilde\sigma_\alpha|^2_{h(\ell)})^{-1}(\Sigma_{\alpha=0}^{N_m}\hat\gamma_\alpha(Y)|\tilde\sigma_\alpha|^2_{h(\ell)}),$$

the left-hand side of (3.10) divided by $m \gg 1$ is written as

$$\int_{M_m}|\mathcal{Y}_{|M_m}|^2_{(\omega_{\rm FS}/m)}\,(\omega_{\rm FS}/m)^n \;=\; \int_M |\bar\partial\phi_Y|^2_{\eta_m}\,\eta_m^n \;\ge\; C_3\int_M \phi_Y^2\,\eta_m^n,$$

for some C_3, because the Kähler manifolds (M,η_m), $m \gg 1$, have bounded geometry (see also Remark 2.11). Hence, by $f_1 < 2f_2$ and (3.10), we see that, for $m \gg 1$,

$$\begin{aligned}
q^n C_1\,\Sigma_{k=1}^{\nu_m}\,n_k b_k^2 \;&\ge\; m\,C_3\int_M \phi_Y^2\,\eta_m^n \;=\; C_3\,q\,f_2\\
&>\; C_3\,q\,f_1/2 \;\ge\; C_4\,q^{n+1}\Sigma_{\alpha=0}^{N_m}\,\hat\gamma_\alpha(Y)^2
\end{aligned}$$

for some C_4, where in the last inequality, we used Remark 2.11. By $\Sigma_{\alpha=0}^{N_m}\gamma_\alpha(Y)=0$, we here observe that $\Sigma_{\alpha=0}^{N_m}\gamma_\alpha(Y)^2 \;\leq\; \Sigma_{\alpha=0}^{N_m}\hat{\gamma}_\alpha(Y)^2$, Hence, whether $f_1\geq 2f_2$ or not, there always exists C_5 such that, for $m\gg 1$,

$$q\,\Sigma_{\alpha=0}^{N_m}\gamma_\alpha(Y)^2 \;\leq\; C_5\,\Sigma_{k=1}^{\nu_m} n_k b_k^2. \tag{3.11}$$

Step 4. Put $P := W+X''$. In view of [26], Theorem 1.6, a weighted version of (3.4.2) in [26] is true (cf. [14], [19]). Hence by $T\in\Delta_1$, we obtain

$$\dot{f}_{X,m}(0) \;=\; \dot{f}_{P,m}(0) \;=\; (n+1)\int_M \frac{\Sigma_{\alpha=0}^{N_m}\gamma_\alpha(P)\,|\tilde{\beta}_\alpha|^2_{h(\ell)}}{\Sigma_{\alpha=0}^{N_m}|\tilde{\beta}_\alpha|^2_{h(\ell)}}\,\Phi_m^*\omega_{\mathrm{FS}}^n, \tag{3.12}$$

where $\gamma_{k,i}(P) := b_k+\gamma_{k,i}(X'')\,(=\gamma_{l(k,i)}(P))$. Let $C_{q,\ell}=1+\Sigma_{j=0}^{\ell}\alpha_j q^{j+1}$ be as in (3.1). Then by (3.1) and (3.3), there exist a function $u_{m,\ell}$ and a 1-form $\theta_{m,\ell}$ on M such that

$$\begin{cases} B_q(\omega(\ell),F(\ell)) \;=\; (n!/m^n)\Sigma_{\alpha=0}^{N_m}|\tilde{\beta}_\alpha|^2_{h(\ell)} \;=\; C_{q,\ell}+u_{m,\ell}\,q^{\ell+2}; \\ \eta_m=(1/m)\Phi_m^*\omega_{\mathrm{FS}}=\omega(\ell)+\theta_{m,\ell}\,q^{\ell+2}, \end{cases} \tag{3.13}$$

where we have the inequalities $\|u_{m,\ell}\|_{C^0(M)}\leq C_6$ and $\|\theta_{m,\ell}\|_{C^0(M,\omega_0)}\leq C_7$ for some C_6 and C_7 (cf. Remark 2.11; see also [25], [14]). Hence, if $m\gg 1$,

$$\frac{|\dot{f}_{X,m}(0)|}{(n+1)!} \;\leq\; \int_M \frac{\left(\Sigma_{\alpha=0}^{N_m}\gamma_\alpha(P)\,|\tilde{\beta}_\alpha|^2_{h(\ell)}\right)\left\{1+\Sigma_{i=1}^{\infty}(-u_{m,\ell}\,q^{\ell+2}/C_{q,\ell})^i\right\}\left\{\omega(\ell)+\theta_{m,\ell}\,q^{\ell+2}\right\}^n}{C_{q,\ell}}.$$

Here, $\{\,1+\Sigma_{i=1}^{\infty}\,(-u_{m,\ell}\,q^{\ell+2}/C_{q,\ell})^i\,\}\{\,\omega(\ell)+\theta_{m,\ell}\,q^{\ell+2}\,\}^n$ is written as $(1+w_{m,\ell})\,\omega(\ell)^n$ for some function $w_{m,\ell}$ on M such that the inequality $\|w_{m,\ell}\|_{C^0(M)}\leq C_8$ holds for some C_8. Then by $\int_M\{\Sigma_{\alpha=0}^{N_m}\gamma_\alpha(P)|\tilde{\beta}_\alpha|^2_{h(\ell)}\}\,\omega(\ell)^n = \Sigma_{k=1}^{\nu_m}\{e^{d\chi_k(F(\ell))}\;\Sigma_{i=1}^{n_k}(b_k+\gamma_{k,i}(X''))\,\}=0$, we have

$$\begin{aligned} |\dot{f}_{X,m}(0)| \;&\leq\; (n+1)!\,q^{\ell+2}\int_M\left|\frac{\Sigma_{\alpha=0}^{N_m}\gamma_\alpha(P)\,|\tilde{\beta}_\alpha|^2_{h(\ell)}}{C_{q,\ell}}\right|\cdot|w_{m,\ell}|\;\omega(\ell)^n \\ &=(n+1)!q^{\ell+2-n}\int_M\{1+(u_{m,\ell}q^{\ell+2}/C_{q,\ell})\}\left|\frac{\Sigma_{\alpha=0}^{N_m}\gamma_\alpha(P)|\tilde{\beta}_\alpha|^2_{h(\ell)}}{\Sigma_{\alpha=0}^{N_m}|\tilde{\beta}_\alpha|^2_{h(\ell)}}\right|\cdot\left|\frac{w_{m,\ell}}{n!}\right|\omega(\ell)^n. \end{aligned}$$

In view of (3.4), by setting $\hat{\phi} := (m\,\Sigma_{\alpha=0}^{N_m}|\tilde{\beta}_\alpha|^2)^{-1}(\Sigma_{\alpha=0}^{N_m}\gamma_\alpha(P)\,|\tilde{\beta}_\alpha|^2)$, there exist C_9 and C_{10} such that, for $m\gg 1$,

$$|\dot{f}_{X,m}(0)| \;\leq\; q^{\ell+1-n}C_9\|\hat{\phi}\|_{L^1(M,\omega(\ell))} \;\leq\; q^{\ell+1-n}C_{10}\|\hat{\phi}\|_{L^2(M,\omega(\ell))}. \tag{3.14}$$

Step 5. Note that $0\leq n_k^2 b_k^2=\{\,\Sigma_{i=1}^{n_k}\gamma_{k,i}(X'')\,\}^2\leq n_k\Sigma_{i=1}^{n_k}\gamma_{k,i}(X'')^2$ holds for all k by the Cauchy-Schwarz inequality. Hence

$$\Sigma_{k=1}^{\nu_m} n_k b_k^2 \;\leq\; \Sigma_{\alpha=0}^{N_m}\gamma_\alpha(X'')^2. \tag{3.15}$$

From $\Sigma_{\alpha=0}^{N_m}\gamma_\alpha(X'')=0$ and $\hat{\gamma}_\alpha(X'')=\gamma_\alpha(X'')+m\,c_{X''}$, it follows that $\Sigma_{\alpha=0}^{N_m}\hat{\gamma}_\alpha(X'')^2 = (N_m+1)(m\,c_{X''})^2+\Sigma_{\alpha=0}^{N_m}\gamma_\alpha(X'')^2$. In particular,

$$\Sigma_{\alpha=0}^{N_m}\gamma_\alpha(X'')^2 \;\leq\; \Sigma_{\alpha=0}^{N_m}\hat{\gamma}_\alpha(X'')^2. \tag{3.16}$$

Since $\gamma_{k,i}(P) = b_k + \gamma_{k,i}(X'')$, (3.15) and (3.16) above imply that

$$\Sigma_{\alpha=0}^{N_m}\gamma_\alpha(P)^2 \;\le\; 2\{\,\Sigma_{k=1}^{\nu_m} n_k b_k^2 \,+\, \Sigma_{\alpha=0}^{N_m}\gamma_\alpha(X'')^2\,\} \;\le\; 4\Sigma_{\alpha=0}^{N_m}\hat{\gamma}_\alpha(X'')^2. \tag{3.17}$$

By $X = Y + P$, we have $\Sigma_{\alpha=0}^{N_m}\gamma_\alpha(X)^2 \le 2\{\,\Sigma_{\alpha=0}^{N_m}\gamma_\alpha(Y)^2 + \Sigma_{\alpha=0}^{N_m}\gamma_\alpha(P)^2\,\}$, because $\gamma_\alpha(X) = \gamma_\alpha(Y) + \gamma_\alpha(P)$. Hence, by (3.11), (3.15), (3.16) and (3.17), we obtain

$$\begin{aligned} q\Sigma_{\alpha=0}^{N_m}\gamma_\alpha(X)^2 \;&\le\; 2q\Sigma_{\alpha=0}^{N_m}\gamma_\alpha(Y)^2 \,+\, 2q\Sigma_{\alpha=0}^{N_m}\gamma_\alpha(P)^2 \\ &\le\; 2C_5\Sigma_{k=1}^{\nu_m} n_k\, b_k^2 \,+\, \Sigma_{\alpha=0}^{N_m}\gamma_\alpha(P)^2 \;\le\; C_{11}\,\Sigma_{\alpha=0}^{N_m}\hat{\gamma}_\alpha(X'')^2 \end{aligned} \tag{3.18}$$

for $m \gg 1$, where we put $C_{11} := 4 + 2C_5$. Fix a positive real number ℓ_0 independent of the choice of m and X. Put $\delta_0 := q^{1/2+\ell_0}/\sqrt{\Sigma_{\alpha=0}^{N_m}\hat{\gamma}_\alpha(X'')^2}$. Then by (3.18), we have $0 < \delta_0 < \sqrt{C_{11}}\,/\bar{\gamma}$, where $\bar{\gamma} := \max\{\,|\gamma_\alpha(X)|\,;\,\alpha = 0, 1, \dots, N_m\,\}$. In view of Step 1 of [15], Section 4, by assuming $|t| \le \delta_0$, we see that the family of Kähler manifolds $(M, q\,\Phi_m^*\lambda_t^*\omega_{\rm FS})$ have bounded geometry.

Step 6. At the beginning of this step, we shall show the inequality (3.19) below as an analogue of [21], (5.9), by proving that an argument of Phong and Sturm [21] for $\dim H = 0$ is valid also for $\dim H > 0$. To see this, we consider the following exact sequence of holomorphic vector bundles

$$0 \to TM_{m,t} \to T\mathbb{P}^*(V_m)_{|M_{m,t}} \to TM_{m,t}^{\perp} \to 0,$$

where $TM_{m,t}^{\perp}$ is regarded as the normal bundle of $M_{m,t}$ in $\mathbb{P}^*(V)$. The pointwise estimate (cf. [21], (5.16)) of the second fundamental form for this exact sequence has nothing to do with $\dim H$, and as in [21], (5.15), it gives the inequality

$$\int_{M_{m,t}} |\mathcal{X}''_{TM_{m,t}^{\perp}}|^2_{\omega_{\rm FS}}\,\omega_{\rm FS}^n \;\ge\; C_{12}\int_{M_{m,t}} |\,\bar{\partial}\mathcal{X}''_{TM_{m,t}^{\perp}}|^2_{\omega_{\rm FS}}\,\omega_{\rm FS}^n.$$

for some C_{12}. Let $\mathcal{A}^{0,p}(T_M)$, $p = 0{,}1$, denote the sheaf of germs of smooth $(0,p)$-forms on M with values in the holomorphic tangent bundle TM of M, and endow M with the Kähler metric $(1/m)\Phi_m^*\lambda_t^*\omega_{\rm FS}$. We then consider the operator $\Box_{TM} := -\bar{\partial}^{\#}\,\bar{\partial}$ on $\mathcal{A}^{0,0}(T_M)$, where $\bar{\partial}^{\#} : \mathcal{A}^{0,1}(T_M) \to \mathcal{A}^{0,0}(T_M)$ is the formal adjoint of $\bar{\partial} : \mathcal{A}^{0,0}(T_M) \to \mathcal{A}^{0,1}(T_M)$. Since by Step 1, the Kähler metrics $q\Phi_m^*\lambda_t^*\omega_{\rm FS}$ has bounded geometry, the first positive eigenvalue λ_1 of the operator $-\Box_{TM}$ on $\mathcal{A}^{0,0}(T_M)$ is bounded from below by C_{13}. Hence, by $X'' \in \mathfrak{p}^{\perp}$,

$$\int_{M_{m,t}} |\bar{\partial}\mathcal{X}''_{TM_{m,t}}|^2_{(\omega_{\rm FS}/m)}\,(\omega_{\rm FS}/m)^n \;\ge\; \lambda_1\int_{M_{m,t}} |\mathcal{X}''_{TM_{m,t}}|^2_{(\omega_{\rm FS}/m)}\,(\omega_{\rm FS}/m)^n.$$

Since $\bar{\partial}\mathcal{X}''_{TM_{m,t}^{\perp}} = -\bar{\partial}\mathcal{X}''_{TM_{m,t}}$, by $\lambda_1 \ge C_{13}$, it now follows that

$$\begin{aligned} \ddot{f}_{X,m}(t) \;&=\; \int_{M_{m,t}} |\mathcal{X}''_{TM_{m,t}^{\perp}}|^2_{\omega_{\rm FS}}\omega_{\rm FS}^n \\ &\ge\; C_{12}C_{13}\,q\int_{M_{m,t}} |\mathcal{X}''_{TM_{m,t}}|^2_{\omega_{\rm FS}}\omega_{\rm FS}^n. \end{aligned} \tag{3.19}$$

In view of the equality $|\mathcal{X}''_{TM_{m,t}}|^2_{\omega_{\rm FS}} + |\mathcal{X}''_{TM^{\perp}_{m,t}}|^2_{\omega_{\rm FS}} = |\mathcal{X}''_{|M_{m,t}}|^2_{\omega_{\rm FS}}$, by adding the integral $C_{12}C_{13}\,q\int_{M_{m,t}} |\mathcal{X}''_{TM^{\perp}_{m,t}}|^2_{\omega_{\rm FS}}\omega^n_{\rm FS}$ to both sides of (3.19) and by dividing the resulting inequality by $(1+C_{12}C_{13}q)$, we see that, for some C_{14} and C_{15},

$$\begin{aligned}\ddot{f}_{X,m}(t) \;&=\; \int_M \Phi_m^*\lambda_t^*\left(|\mathcal{X}''_{TM^{\perp}_{m,t}}|^2_{\omega_{\rm FS}}\,\omega^n_{\rm FS}\right)\\ &\geq\; C_{14}\,q\int_M \Phi_m^*\lambda_t^*\left(|\mathcal{X}''_{|M_{m,t}}|^2_{\omega_{\rm FS}}\,\omega^n_{\rm FS}\right)\\ &\geq\; C_{15}\,q\int_{M_m} |\mathcal{X}''_{|M_m}|^2_{\omega_{\rm FS}}\,\omega^n_{\rm FS} \;\geq\; C_{15}\,q\int_M \Theta\,\Phi_m^*\omega^n_{\rm FS},\end{aligned}\tag{3.20}$$

where $\Theta := (\Sigma_{\alpha=0}^{N_m}|\tilde{\beta}_\alpha|^2)^{-2}\{\,(\Sigma_{\alpha=0}^{N_m}|\tilde{\beta}_\alpha|^2)(\Sigma_{\alpha=0}^{N_m}\hat{\gamma}_\alpha(X'')^2|\tilde{\beta}_\alpha|^2) - (\Sigma_{\alpha=0}^{N_m}\hat{\gamma}_\alpha(X'')\,|\tilde{\beta}_\alpha|^2)^2\,\}$ is nonnegative everywhere on M. Then by (3.14) and (3.20),

$$\begin{cases}\dot{f}_{X,m}(\delta_0) & \geq\; \dot{f}_{X,m}(0) + C_{15}\,\delta_0\,q\int_M \Theta\Phi_m^*\omega^n_{\rm FS}\\ & \geq\; -q^{\ell+1-n}C_{10}\|\hat{\phi}\|_{L^2(M,\omega(\ell))} + C_{15}\,\delta_0\,q\int_M \Theta\Phi_m^*\omega^n_{\rm FS},\\ \dot{f}_{X,m}(-\delta_0) & \leq\; \dot{f}_{X,m}(0) - C_{15}\,\delta_0\,q\int_M \Theta\Phi_m^*\omega^n_{\rm FS}\\ & \leq\; q^{\ell+1-n}C_{10}\|\hat{\phi}\|_{L^2(M,\omega(\ell))} - C_{15}\,\delta_0\,q\int_M \Theta\Phi_m^*\omega^n_{\rm FS}.\end{cases}\tag{3.21}$$

By (3.20) and [15], Lemma 3.4, the proof of Main Theorem is reduced to showing the following three conditions for all $m \gg 1$:

$$\text{i) } \dot{f}_{X,m}(\delta_0) > 0 > \dot{f}_{X,m}(-\delta_0), \quad \text{ii) } \int_M \Theta\Phi_m^*\omega^n_{\rm FS} > 0, \quad \text{iii) } t_X^{(m)}\cdot X \in U_m.$$

Since iii) follows from Remark 3.31 below, we have only to prove i) and ii). Then by (3.21), it suffices to show the following for all $m \gg 1$:

$$C_{15}\,\delta_0\,q\int_M \Theta\Phi_m^*\omega^n_{\rm FS} \;-\; C_{10}\,q^{\ell+1-n}\|\hat{\phi}\|_{L^2(M,\omega(\ell))} \;>\; 0.\tag{3.22}$$

Let us define real numbers $\hat{e}_1$, $\hat{e}_2$, e_1, e_2 by setting

$$\begin{aligned}\hat{e}_1 &:= \int_M \frac{\Sigma_{\alpha=0}^{N_m}\hat{\gamma}_\alpha(X'')^2|\tilde{\beta}_\alpha|^2}{\Sigma_{\alpha=0}^{N_m}|\tilde{\beta}_\alpha|^2}\,\omega(\ell)^n, & \hat{e}_2 &:= \int_M \left(\frac{\Sigma_{\alpha=0}^{N_m}\hat{\gamma}_\alpha(X'')|\tilde{\beta}_\alpha|^2}{\Sigma_{\alpha=0}^{N_m}|\tilde{\beta}_\alpha|^2}\right)^2\omega(\ell)^n,\\ e_1 &:= \int_M \frac{\Sigma_{\alpha=0}^{N_m}\gamma_\alpha(P)^2|\tilde{\beta}_\alpha|^2}{\Sigma_{\alpha=0}^{N_m}|\tilde{\beta}_\alpha|^2}\omega(\ell)^n, & e_2 &:= \int_M \left(\frac{\Sigma_{\alpha=0}^{N_m}\gamma_\alpha(P)|\tilde{\beta}_\alpha|^2}{\Sigma_{\alpha=0}^{N_m}|\tilde{\beta}_\alpha|^2}\right)^2\omega(\ell)^n.\end{aligned}$$

By the Cauchy-Schwarz inequality, we always have $\hat{e}_1 \geq \hat{e}_2$ and $e_1 \geq e_2$. Now, the following cases are possible:

Case 1: $\hat{e}_1 > 2\,\hat{e}_2$, Case 2: $\hat{e}_1 \leq 2\,\hat{e}_2$.

In view of the identities in (3.13), we can write

$$\begin{aligned} \hat{e}_1 \;&=\; q^n n! \int_M \frac{\Sigma_{\alpha=0}^{N_m} \hat{\gamma}_\alpha(X'')^2 |\tilde{\beta}_\alpha|^2_{h(\ell)}}{1+\Sigma_{\alpha=0}^{\ell}\alpha_k q^{k+1} + u_{m,\ell}\, q^{\ell+2}}\,\omega(\ell)^n,\\ e_1 \;&=\; q^n n! \int_M \frac{\Sigma_{\alpha=0}^{N_m} \gamma_\alpha(P)^2 |\tilde{\beta}_\alpha|^2_{h(\ell)}}{1+\Sigma_{\alpha=0}^{\ell}\alpha_k q^{k+1} + u_{m,\ell}\, q^{\ell+2}}\,\omega(\ell)^n,\\ \int_M \Theta\Phi_m^*\omega_{\mathrm{FS}}^n \;&=\; m^n \int_M \Theta\{\omega(\ell) + \theta_{m,\ell}\, q^{\ell+2}\}^n, \end{aligned}$$

and hence, given a positive real number $0 < \varepsilon \ll 1$, both $\hat{e}_1$ and $\int_M \Theta\Phi_m^*\omega_{\mathrm{FS}}^n$ above are estimated, for all $m \gg 1$, by

$$(1-\varepsilon)\, q^n\, \{\Sigma_{\alpha=0}^{N_m}\hat{\gamma}_\alpha(X'')^2\} \;\le\; \hat{e}_1/n! \;\le (1+\varepsilon)\, q^n\, \{\Sigma_{\alpha=0}^{N_m}\hat{\gamma}_\alpha(X'')^2\}, \tag{3.23}$$

$$(1-\varepsilon)\, q^n\, \{\Sigma_{\alpha=0}^{N_m}\gamma_\alpha(P)^2\} \;\le\; e_1/n! \;\le (1+\varepsilon)\, q^n\, \{\Sigma_{\alpha=0}^{N_m}\gamma_\alpha(P)^2\}, \tag{3.24}$$

$$(1-\varepsilon)\, q^{-n}\int_M \Theta\omega(\ell)^n \;\le\; \int_M \Theta\Phi_m^*\omega_{\mathrm{FS}}^n \;\le\; (1+\varepsilon)\, q^{-n}\int_M \Theta\omega(\ell)^n, \tag{3.25}$$

where we used Remark 2.11. Moreover, we can write e_2 in the form

$$q^{-2}\|\hat{\phi}\|^2_{L^2(M,\omega(\ell))} \;=\; e_2 \;\le\; e_1. \tag{3.26}$$

Step 7. We first consider Case 1. Then from (3.17), (3.23), (3.24), (3.25), (3.26), $\hat{e}_2 < \hat{e}_1/2$ and the definition of δ_0, it follows that

$$\begin{aligned} &\text{L.H.S. of (3.22)} \;\ge\; (1-\varepsilon)\, C_{15}\, \delta_0\, q^{1-n}\int_M \Theta\omega(\ell)^n \;-\; q^{\ell+1-n} C_{10}\|\hat{\phi}\|_{L^2(M,\omega(\ell))}\\ &\ge\; (1-\varepsilon)\, C_{15}\, q^{1-n}\, \delta_0\, (\hat{e}_1 - \hat{e}_2) \;-\; q^{\ell+2-n} C_{10}\sqrt{e_1}\\ &\ge\; (1-\varepsilon)\, C_{15}\, \delta_0\, q^{1-n}\, \hat{e}_1/2 - \; (1+\varepsilon)^{1/2} C_{10}\, q^{\ell+2-\frac{n}{2}}\{\, n!\Sigma_{\alpha=0}^{N_m}\gamma_\alpha(P)^2\,\}^{1/2}\\ &\ge\; (1-\varepsilon)^2 C_{15}\, \delta_0\, q\,\{\,\Sigma_{\alpha=0}^{N_m}\hat{\gamma}_\alpha(X'')^2\,\}\, n!/2\\ &\qquad\qquad -\; 2(1+\varepsilon)^{1/2} C_{10}\, q^{\ell+2-\frac{n}{2}}\{\, n!\,\Sigma_{\alpha=0}^{N_m}\hat{\gamma}_\alpha(X'')^2\,\}^{1/2}\\ &\ge\; \{n!\,\Sigma_{\alpha=0}^{N_m}\hat{\gamma}_\alpha(X'')^2\,\}^{1/2}\left\{\sqrt{n!}\,(1-\varepsilon)^2 C_{15} q^{\ell_0+\frac{3}{2}}/2 - 2(1+\varepsilon)^{1/2} C_{10} q^{\ell+2-\frac{n}{2}}\right\}, \end{aligned}$$

for $m \gg 1$. Now we see that, if $\ell > (n-1)/2 + \ell_0$, then $q^{\ell+2-\frac{n}{2}}/q^{\ell_0+\frac{3}{2}}$ converges to 0 as $m \to \infty$. Thus, if $m \gg 1$, then by choosing ℓ such that $\ell > (n-1)/2 + \ell_0$, we now see from the computation above that L.H.S. of (3.22) is positive, as required.

Step 8. Let us finally consider Case 2. For each fixed ℓ, the Kähler form η_m converges to ω_0, as $m \to \infty$, in $C^j(M)$-norm for all positive integers j (cf. [25], [14]; see also Remark 2.11). Note that $X'' \in \mathfrak{p}^\perp$. In view of $\int_M \phi_{X''}\eta_m^n = 0$, we see that

$$\|\phi_{X''}\|^2_{L^2(M,\eta_m)} \;\le\; C_{16}\, \|\bar{\partial}\phi_{X''}\|^2_{L^2(M,\eta_m)} \;=\; C_{16}\, \|\Phi_m^*\mathcal{X}''_{TM_m}\|^2_{L^2(M,\eta_m)}, \tag{3.27}$$

for some C_{16}, where by abuse of terminology, the differential $(\Phi_m^{-1})_* : TM_m \to TM$ is denoted by Φ_m^*. Moreover, by (3.19) applied to $t = 0$, we obtain

$$\|\,\Phi_m^*\mathcal{X}''_{TM_m}\|^2_{L^2(M,\eta_m)} \;\le\; (C_{12}C_{13})^{-1} q^{-1}\|\Phi_m^*\mathcal{X}''_{TM_m^\perp}\|^2_{L^2(M,\eta_m)}. \tag{3.28}$$

From now on until the end of this proof, we assume that $m \gg 1$. By (3.27) together with (3.28) and (3.7), there exist C_{17} and C_{18} such that

$$(3.29) \qquad \begin{cases} \|\Phi_m^* \mathcal{X}''_{TM_m^\perp}\|_{L^2(M,\eta_m)} \;\geq\; C_{17}\, q^{1/2}\, \|\phi_{X''}\|_{L^2(M,\eta_m)} \\ \geq\; C_{18}\, q^{1/2}\, \|\phi_{X''}\|_{L^2(M,\omega(\ell))} \;=\; C_{18}\, q^{3/2}\, \sqrt{\hat{e}_2}. \end{cases}$$

We now observe the pointwise estimate $q^{1/2}|\mathcal{X}''_{|M_m}|_{\omega_{\mathrm{FS}}} = |\mathcal{X}''_{|M_m}|_{\eta_m} \geq |\mathcal{X}''_{TM_m^\perp}|_{\eta_m}$. Hence by (3.20) and (3.29), we obtain

$$(3.30) \qquad \begin{cases} \ddot{f}_{X,m}(t) \;\geq\; C_{15} \int_{M_m} \left(q^{1/2}|\mathcal{X}''_{|M_m}|_{\omega_{\mathrm{FS}}}\right)^2 \omega_{\mathrm{FS}}^n \\ \geq\; C_{15}\, q^{-n} \|\Phi_m^* \mathcal{X}''_{TM_m^\perp}\|^2_{L^2(M,\eta_m)} \;\geq\; C_{19}\, q^{3-n}\hat{e}_2, \end{cases}$$

for some C_{19}. As in deducing (3.21) from (3.14) and (3.20), we obtain by (3.14) and (3.30) the inequalities

$$\dot{f}_{X,m}(\delta_0) \;\geq\; R \quad \text{and} \quad \dot{f}_{X,m}(-\delta_0) \;\leq\; -R,$$

where $R := -q^{\ell+1-n} C_{10}\|\hat{\phi}\|_{L^2(M,\omega(\ell))} + C_{19}\,\delta_0\, q^{3-n}\hat{e}_2$. Hence, it suffices to show that $R > 0$. In view of the definition of $\hat{\phi}$ and e_2, we see from $\hat{e}_1 \leq 2\hat{e}_2$ and (3.26) that

$$R \;=\; C_{19}\,\delta_0\, q^{3-n}\hat{e}_2 \;-\; C_{10}\, q^{\ell+2-n}\sqrt{e_2} \;\geq\; C_{19}\,\delta_0\, q^{3-n}\hat{e}_1/2 \;-\; C_{10}\, q^{\ell+2-n}\sqrt{e_1}.$$

Here by (3.23) and (3.24), we obtain

$$\begin{aligned} \frac{\delta_0 q^{3-n}\hat{e}_1}{q^{\ell+2-n}\sqrt{e_1}} \;&=\; q^{3/2+\ell_0-\ell} \sqrt{\frac{\hat{e}_1}{e_1}} \sqrt{\frac{\hat{e}_1}{\Sigma_{\alpha=0}^{N_m}\hat{\gamma}_\alpha(X'')^2}} \\ &\geq\; C_{20}\, q^{(3+n)/2+\ell_0-\ell} \sqrt{\frac{\Sigma_{\alpha=0}^{N_m}\,\hat{\gamma}_\alpha(X'')^2}{\Sigma_{\alpha=0}^{N_m}\,\gamma_\alpha(P)^2}} \;\geq\; \frac{C_{20}}{2}\, q^{(3+n)/2+\ell_0-\ell} \end{aligned}$$

for some C_{20}, where the last inequality follows from (3.17). Therefore, by choosing ℓ such that $\ell > (3+n)/2 + \ell_0$, we now conclude that $R > 0$ for $m \gg 1$, as required. □

Remark 3.31. In the above proof, it is easy to check the condition iii) in Step 6 as follows: In view of $|t_X^{(m)}| < \delta_0$, it suffices to show that, if $m \gg 1$, then

$$(3.32) \qquad t \cdot X \in U_m, \qquad \text{for all } (t, X) \in \mathbb{R} \times \mathfrak{p}''_m \text{ with } |t| < \delta_0.$$

For each $Q \in \mathfrak{p}$, let $u_Q \in C^\infty(M)_{\mathbb{R}}$ denote the Hamiltonian function for the holomorphic vector field Q on the Kähler manifold (M, ω_0) characterized by the equalities

$$i_Q\omega_0 \;=\; (\sqrt{-1}/2\pi)\,\bar{\partial} u_Q \quad \text{and} \quad \int_M u_Q \omega_0^n \;=\; 0.$$

Define compact subsets $\Sigma(\mathfrak{p}'_m)$, $\Sigma(\mathfrak{p})$ of $\mathfrak{p}$ by setting

$$\begin{cases} \Sigma(\mathfrak{p}'_m) & := \{\, Q \in \mathfrak{p}'_m \,;\, \|\bar{\partial} u_Q\|_{L^2(M,\omega_0)} = 1 \,\}, \\ \Sigma(\mathfrak{p}) & := \{\, Q \in \mathfrak{p} \,;\, \|\bar{\partial} u_Q\|_{L^2(M,\omega_0)} = 1 \,\}. \end{cases}$$

Choose an orthonormal basis $\mathcal{S} := \{s_0, s_1, \dots, s_{N_m}\}$ for the Hermitian vector space $(V_m, \rho_{h(\ell)})$. For the space $\mathcal{H}_m$ of all Hermitian matrices of order $N_m + 1$, define a norm

$$\mathcal{H}_m \to \mathbb{R}_{\geq 0}, \qquad A = (a_{\alpha\beta}) \;\mapsto\; \|A\|_m := \sqrt{\operatorname{tr} A^* A} = \sqrt{\Sigma_{\alpha,\beta}\, |a_{\alpha\beta}|^2}$$

on $\mathcal{H}_m$. Let $m \gg 1$. The infinitesimal action of $\mathfrak{p}_m$ on V_m is given by

$$Q \cdot s_\beta \;=\; \Sigma_{\alpha=0}^{N_m}\, s_\alpha\, \gamma_{\alpha\beta}(Q), \qquad\qquad Q \in \mathfrak{p}_m,$$

where $\gamma_Q = (\gamma_{\alpha\beta}(Q)) \in \mathcal{H}_m$ denotes the representation matrix of Q on V_m with respect to $\mathcal{S}$. Let $X \in \mathfrak{p}''_m$, and let δ_0 be as in Step 5 above. For $t \in \mathbb{R}$ with $|t| < \delta_0$, we put $\tilde{X} := t\,X$. In order to prove (3.32) above, it suffices to show

$$\zeta(\operatorname{ad}\tilde{X})Q \notin \mathfrak{p}''_m \qquad \text{for all } Q \in \Sigma(\mathfrak{p}'_m). \tag{3.33}$$

Let $Q \in \Sigma(\mathfrak{p}'_m)$. For a suitable choice of a basis $\mathcal{S}$ as above, we may assume that the representation matrix γ_Q of Q is a real diagonal matrix. Note also that $\operatorname{tr}\gamma_Q = 0$. Let $\Phi_m : M \to \mathbb{P}^{N_m}(\mathbb{C})$ be the Kodaira embedding of M defined by (cf. (3.2))

$$\Phi_m(p) := (\tilde{s}_0(p) : \tilde{s}_1(p) : \cdots : \tilde{s}_{N_m}(p)).$$

In view of the definition $\eta_m := \Phi_m^*\omega_{\mathrm{FS}}/m$ of η_m, the Hamiltonian function ϕ_Q on (M, η_m) associated to the holomorphic vector field Q is expressed in the form

$$\phi_Q \;=\; (\Sigma_{\alpha=0}^{N_m}\,\hat{\gamma}_{\alpha\alpha}(Q)\,|\tilde{s}_\alpha|^2)/(m\Sigma_{\alpha=0}^{N_m}\,|\tilde{s}_\alpha|^2).$$

We define $\hat{\gamma}_Q := (\hat{\gamma}_{\alpha\beta}(Q)) \in \mathcal{H}_m$ by setting $\hat{\gamma}_{\alpha\beta}(Q) := \{\gamma_{\alpha\alpha}(Q) + m\,c_Q\}\delta_{\alpha\beta}$ for Kronecker's delta $\delta_{\alpha\beta}$. As in deducing (3.16) from $\hat{\gamma}_\alpha(X'') = \gamma_\alpha(X'') \,+\, m\,c_{X''}$, we easily see that

$$\|\gamma_Q\|_m^2 \;\le\; \|\hat{\gamma}_Q\|_m^2.$$

Recall that η_m is expressible as $\omega_0 + (\sqrt{-1}/2\pi)\,q\,\partial\bar{\partial}\xi_m$ for some real-valued smooth function ξ_m on M such that

$$\|\xi_m\|_{C^3(M)} \le C_{21}, \tag{3.34}$$

where all C_j's in this remark are positive constants independent of the choice of m, X and Q. We now observe that

$$\phi_Q \;=\; u_Q \,+\, q\,(Q\xi_m). \tag{3.35}$$

Note that $Q \in \Sigma(\mathfrak{p}'_m) \subset \Sigma(\mathfrak{p})$. Since Q sits in the compact set $\Sigma(\mathfrak{p})$, and since $\Sigma(\mathfrak{p})$ is independent of the choice of m, there exist C_{22} and C_{23} such that

$$0 \;<\; C_{22} \;\le\; \int_M u_Q^2\omega_0^n \left(= \int_M \phi_Q^2\eta_m^n\right) \le\; C_{23}.$$

Note that both η_m and $\omega(\ell)$ converge to ω_0 as $m \to \infty$ (see (3.3) and the statement at the beginning of Step 8). Note also that, by Remark 2.11, the function $(n!/m^n)\Sigma_{\alpha=0}^{N_m}\,|\tilde{s}_\alpha|^2_{h(\ell)}$ on M converges uniformly to 1, as $m \to \infty$. Again by Remark 2.11, it now follows from the Cauchy-Schwarz inequality that, for $m \gg 1$,

$$\begin{aligned} \|\gamma_Q\|_m^2 \;&=\; \sum_{\alpha=0}^{N_m}\gamma_{\alpha\alpha}(Q)^2 \;\ge\; C_{24}\,m^n \int_M \frac{\Sigma_{\alpha=0}^{N_m}\,\gamma_{\alpha\alpha}(Q)^2\,|\tilde{s}_\alpha|^2}{\Sigma_{\alpha=0}^{N_m}\,|\tilde{s}_\alpha|^2}\,\omega(\ell)^n \\ &\ge\; C_{24}\,m^{n+2}\int_M (\Phi_m^*\varphi_Q)^2\,\omega(\ell)^n \;\ge\; C_{25}\,m^{n+2}\int_M (\Phi_m^*\varphi_Q)^2\,\eta_m^n \end{aligned} \tag{3.36}$$

for some C_{24} and C_{25}, where $\Phi_m^*\varphi_Q$ is as in (3.4). Then for $m \gg 1$,

$$\begin{aligned} C_{26} \;&=\; \max_{J\in\Sigma(\mathfrak{p})} \int_M |J|^2_{\omega_0}\,\omega_0^n \;\geq\; \int_M |Q|^2_{\omega_0}\,\omega_0^n \;\geq\; C_{27}\int_M |Q|^2_{\eta_m}\,\eta_m^n \\ &=\; C_{27}\,m \int_M \left\{ \frac{\Sigma_{\alpha=0}^{N_m}\,\hat{\gamma}_{\alpha\alpha}(Q)^2\,|\tilde{s}_\alpha|^2_{h(\ell)}}{m^2\Sigma_{\alpha=0}^{N_m}\,|\tilde{s}_\alpha|^2_{h(\ell)}} \;-\; \phi_Q^2 \right\} \eta_m^n \\ &\geq\; \frac{C_{28}}{m^{n+1}} \left\{ \int_M \Sigma_{\alpha=0}^{N_m}\,\hat{\gamma}_{\alpha\alpha}(Q)^2\,|\tilde{s}_\alpha|^2_{h(\ell)}\,\eta_m^n \right\} \;-\; C_{29}\,m \\ &\geq\; \frac{C_{30}}{m^{n+1}} \left\{ \int_M \Sigma_{\alpha=0}^{N_m}\,\hat{\gamma}_{\alpha\alpha}(Q)^2\,|\tilde{s}_\alpha|^2_{h(\ell)}\,\omega(\ell)^n \right\} \;-\; C_{29}\,m \\ &\geq\; C_{31}\frac{\|\hat{\gamma}_Q\|^2_m}{m^{n+1}} \;-\; C_{29}\,m, \end{aligned}$$

for some C_{26}, C_{27}, C_{28}, C_{29}, C_{30} and C_{31}. Hence, if $m \gg 1$, then

$$\|\gamma_Q\|^2_m \;\leq\; \|\hat{\gamma}_Q\|^2_m \;\leq\; C_{32}\,m^{n+2}. \tag{3.37}$$

for some C_{32}. Now by $\zeta(0) = 1$, we define a real analytic function $\tilde{\zeta} = \tilde{\zeta}(x)$ on $\mathbb{R}$ satisfying $\tilde{\zeta}(0) = 0$ by

$$\tilde{\zeta}(x) := \zeta(x) - 1.$$

For $X \in \mathfrak{p}''_m$ above, by choosing an orthonormal basis for $(V_m, \rho_{h(\ell)})$ possibly distinct from the original one, we may assume that the representation matrix γ_X of X is a real diagonal matrix. Recall that $\tilde{X} = t\,X$, where $|t| < \delta_0 := q^{1/2+\ell_0}/\|\hat{\gamma}_{X''}\|_m$. Put $\tilde{X}'' := t\,X''$. Then by (3.18),

$$\sqrt{q/C_{11}}\,\|\gamma_{\tilde{X}}\|_m \;\leq\; \|\hat{\gamma}_{\tilde{X}''}\|_m \;=\; |t|\cdot\|\hat{\gamma}_{X''}\|_m \;\leq\; q^{1/2+\ell_0},$$

i.e., $\|\gamma_{\tilde{X}}\|_m \;\leq\; \sqrt{C_{11}}\,q^{\ell_0}$. Hence, if $m \gg 1$,

$$\|\,\gamma_{\tilde{\zeta}(\operatorname{ad}\tilde{X})Q}\,\|_m \;\leq\; C_{33}\,q^{\ell_0}\,\|\gamma_Q\|_m, \tag{3.38}$$

for some C_{33}. Now by the same argument as in (3.36), we see that, for some C_{34},

$$\|\gamma_{\tilde{\zeta}(\operatorname{ad}\tilde{X})Q}\,\|^2_m \;\geq\; C_{34}\,m^{n+2}\int_M \Phi_m^*\varphi^2_{\tilde{\zeta}(\operatorname{ad}\tilde{X})Q}\,\eta_m^n, \qquad \text{if } m \gg 1. \tag{3.39}$$

Put $a_m := \sqrt{\int_M \Phi_m^*\varphi^2_{\tilde{\zeta}(\operatorname{ad}\tilde{X})Q}\eta_m^n}$. Then for $m \gg 1$, by (3.37), (3.38) and (3.39),

$$a_m \;\leq\; C_{35}\,q^{\ell_0} \tag{3.40}$$

for some C_{35}. Consider the Laplacians $\Box_{\eta_m}$ and $\Box_{\omega_0}$ on functions for the Kähler manifolds (M, η_m) and (M, ω_0), respectively. Note that $\zeta(\operatorname{ad}\tilde{X})Q = Q + \tilde{\zeta}(\operatorname{ad}\tilde{X})Q$. Then for $m \gg 1$, by (3.35), we obtain

$$\left\{ \begin{aligned} &\left|\textstyle\int_M(\Box_{\eta_m}\phi_Q)\phi_{\zeta(\operatorname{ad}\tilde{X})Q}\eta_m^n\right| \\ &=\; \left|\textstyle\int_M(\Box_{\eta_m}\phi_Q)\left(\phi_Q + \Phi_m^*\varphi_{\tilde{\zeta}(\operatorname{ad}\tilde{X})Q}\right)\,\eta_m^n\right| \\ &\geq\; \|\bar{\partial}\phi_Q\|^2_{L^2(M,\eta_m)} \;-\; \left|\textstyle\int_M(\Box_{\eta_m}\phi_Q)(\Phi_m^*\varphi_{\tilde{\zeta}(\operatorname{ad}\tilde{X})Q})\,\eta_m^n\right| \\ &\geq\; \|\bar{\partial}\{u_Q + q(Q\xi_m)\}\|^2_{L^2(M,\eta_m)} \\ &\qquad\qquad -a_m\|\Box_{\eta_m}\{u_Q + q(Q\xi_m)\}\|_{L^2(M,\eta_m)} \\ &\geq\; (1-\epsilon)\,R_m \;-\; (1+\epsilon)\,a_m S_m, \end{aligned} \right. \tag{3.41}$$

where we put $R_m := \|\bar\partial\{u_Q + q(Q\xi_m)\}\|^2_{L^2(M,\omega_0)}$ and $S_m := \|\Box_{\omega_0}\{u_Q + q(Q\xi_m)\}\|_{L^2(M,\omega_0)}$, and $\epsilon \ll 1$ is a positive constant independent of the choice of m, X and Q. Since Q belongs to the compact set $\Sigma(\mathfrak{p})$, by (3.34) and the equality $\|\bar\partial u_Q\|_{L^2(M,\omega_0)} = 1$, we obtain constants C_{36} and C_{37} such that

$$\left\{ \begin{array}{l} R_m \;\geq 1 \,-\, 2\,q\,\|\bar\partial(Q\xi_m)\|_{L^2(M,\omega_0)} \;\geq\; 1 \,-\, C_{36}\,q, \\ S_m \;\leq\; \|\Box_{\omega_0} u_Q\|_{L^2(M,\omega_0)} \,+\, q\,\|\Box_{\omega_0}(Q\xi_m)\|_{L^2(M,\omega_0)} \;\leq\; C_{37}. \end{array} \right. \tag{3.42}$$

Then for $m \gg 1$, by (3.40), (3.41) and (3.42), we finally obtain

$$\left| \int_M (\Box_{\eta_m}\phi_Q)\,\phi_{\zeta(\mathrm{ad}\,\tilde X)Q}\,\eta_m^n \right| \;\geq\; (1-\epsilon)\,(1 - C_{36}q) \,-\, (1+\epsilon)\,C_{35}C_{37}\,q^{\ell_0} \;>\; 0,$$

which implies (3.33), as required.

References

[1] E. Calabi : *Extremal Kähler metrics II*, in "Differential Geometry and Complex Analysis" (ed. I. Chavel, H. M. Farkas), Springer-Verlag, Heidelberg, 1985, 95–114, MR0780039, Zbl0574.58006.

[2] D. Catlin : *The Bergman kernel and a theorem of Tian*, in "Analysis and Geometry in Several Complex Variables" (ed. G. Komatsu, M. Kuranishi), Trends in Math., Birkhäuser, Boston, 1999, 1–23, MR1699887, Zbl0941.32002.

[3] S. K. Donaldson: *Scalar curvature and projective embeddings, I*, J. Differential Geom. **59** (2001), 479–522, MR1916953, Zblpre02053206.

[4] A. Fujiki : *Moduli space of polarized algebraic manifolds and Kähler metrics*, Sugaku **42** (1990), 231–243; English translation: Sugaku Expositions **5** (1992), 173–191, MR1207204, Zbl0796.32009.

[5] A. Futaki and T. Mabuchi : *Moment maps and symmetric multilinear forms associated with symplectic classes*, Asian J. Math. **6** (2002), 349–372, MR1928634, Zblpre01925763.

[6] D. Gieseker: *Global moduli for surfaces of general type*, Invent. Math. **43** (1977), 233–282, MR0498596 Zbl0389.14006.

[7] S. Helgason : *Differential Geometry and Symmetric Spaces*, Pure and Appl. Math. **12**, Academic Press, New York, 1962, MR0145455, Zbl0111.18101.

[8] S. Kobayashi : *Transformation groups in differential geometry*, Springer-Verlag, Berlin-Heidelberg-New York, 1972, MR0355886, Zbl0246.53031.

[9] A. Lichnérowicz: *Isométrie et transformations analytique d'une variété kählérienne compacte*, Bull. Soc. Math. France **87** (1959), 427–437, MR0114187, Zbl0192.28403.

[10] Z. Lu: On the lower order terms of the asymptotic expansion of Tian-Yau-Zelditch, Amer. J. Math. **122** (2000), 235–273, MR1749048, Zbl0972.53042.

[11] M. Lübke and A. Teleman: *The Kobayashi-Hitchin correspondence*, World-Scientific, Singapore, 1995, MR1370660, Zbl0849.32020.

[12] T. Mabuchi : *The Hitchin-Kobayashi correspondence for vector bundles and manifolds*, Japanese, Proc. 48th Geometry Symposium, Ibaraki, August, 2001, 461–468.

[13] T. Mabuchi : *An obstruction to asymptotic semistability and approximate critical metrics*, arXiv: math. DG/0404210, to appear in Osaka J. Math. **41** (2004), 1–20.

[14] T. Mabuchi : *Stability of extremal Kähler manifolds*, arXiv: math. DG/0404211, to appear in Osaka J. Math. **41** (2004), 463–472.

[15] T. Mabuchi : *An energy-theoretic approach to the Hitchin-Kobayashi correspondence for manifolds, I*, to appear in Invent. Math.

[16] T. Mabuchi : *Uniqueness of extremal Kähler metrics for an integral Kähler class*, to appear in Internat. J. Math.

[17] T. Mabuchi and Y. Nakagawa: *Addendum to "The Bando-Calabi-Futaki Character as an Obstruction to Semistability"*, preprint.

[18] D. Mumford : *Varieties defined by quadratic equations*, Questions on Algebraic Varieties (C.I.M.E., 3. Ciclo, Varenna, 1969), Edizioni Cremonese, Rome, 1970, 29–100, MR0282975, Zbl0198.25801.

[19] T. Mabuchi and L. Weng : *Kähler-Einstein metrics and Chow-Mumford stability*, 1998, preprint.

[20] D. Mumford, J. Fogarty and F. Kirwan: Geometric invariant theory, 3rd edition, Ergebnisse der Math. und ihrer Grenzgebiete **34**, Springer-Verlag, 1994, 1–292, MR1304906, Zbl0797.14004.

[21] D. H. Phong and J. Sturm: *Scalar curvature, moment maps, and the Deligne pairing*, arXiv: math. DG/0209098.

[22] G. Tian: *On a set of polarized Kähler metrics on algebraic manifolds*, J. Differential Geom. **32** (1990), 99–130, MR1064867, Zbl0706.53036.

[23] G. Tian : *Kähler-Einstein metrics with positive scalar curvature*, Invent. Math. **130** (1997), 1–37, MR1471884, Zbl0892.53027.

[24] E. Viehweg: *Quasi-projective moduli for polarized manifolds*, Ergebnisse der Math. und ihrer Grenzgebiete, 3. Folge. **30**, Springer-Verlag, 1995, 1–320, MR1368632, Zbl0844.14004.

[25] S. Zelditch : *Szegö kernels and a theorem of Tian*, Internat. Math. Res. Notices **6** (1998), 317–331, MR1616718, Zbl0922.58082.

[26] S. Zhang : *Heights and reductions of semi-stable varieties*, Compositio Math. **104** (1996), 77–105, MR1420712, Zbl0924.11055.

Department of Mathematics, Graduate School of Science, Osaka University, Toyonaka, Osaka, 560-0043 Japan